

DISTANT RED GALAXIES IN THE HUBBLE ULTRA DEEP FIELD¹

S. TOFT², P. VAN DOKKUM², M. FRANX³, R. I. THOMPSON⁴, G. D. ILLINGWORTH⁵, R. J. BOUWENS⁵, M. KRIEK³
Draft version February 2, 2008

ABSTRACT

We take advantage of the Hubble Ultra Deep Field (UDF) data to study the restframe optical and ultra violet (UV) morphologies of the novel population of Distant Red Galaxies (DRGs). Six galaxies with $J - K_s > 2.3$ are found to $K_s = 21.5$, five of which have photometric redshifts $z_{phot} \gtrsim 2$, corresponding to a surface density of 0.9 arcmin^{-2} . The surface brightness distributions of the $z_{phot} \gtrsim 2$ galaxies are better represented by exponential disks than $R^{1/4}$ -laws. Two of the $z_{phot} \gtrsim 2$ galaxies are extended, while three have compact morphologies. The restframe optical morphology of the $z_{phot} \gtrsim 2$ galaxies is quite different from the restframe UV morphology: all the galaxies have red central components which dominate in the NICMOS H_{160} -band images, and distinct off-center blue features which show up in (and often dominate) the ACS images. The mean measured effective radius of the $z_{phot} \gtrsim 2$ galaxies is $\langle r_e \rangle = 1.9 \pm 1.4 \text{ kpc}$, similar (within the errors) to the mean size of LBGs at similar redshifts. All the DRGs are resolved in the ACS images, while four are resolved in the NICMOS images. Two of the $z_{phot} \gtrsim 2$ galaxies are bright X-ray sources and hence host AGN. One of these galaxies is resolved in the ACS and NICMOS images, which means the AGN does not dominate its restframe UV-optical SED, while the other is unresolved in the NICMOS images and hence could have an AGN dominated SED. The diverse restframe optical and UV morphological properties of DRGs derived here suggest that they have complex stellar populations, consisting of both evolved populations that dominate the mass and the restframe optical light, and younger populations, which show up as patches of star formation in the restframe UV light; in many ways resembling the properties of normal local galaxies. This interpretation is supported by fits to the broadband SEDs, which for all five $z_{phot} \gtrsim 2$ are best represented by models with extended star formation histories and substantial amounts of dust.

Subject headings: cosmology: observations — galaxies: high-redshift — galaxies: evolution — galaxies: formation

1. INTRODUCTION

The Faint InfraRed ExtraGalactic Survey (FIRES, Labbé et al. 2003) discovered a highly clustered (Daddi et al. 2003) population of Distant ($z_{phot} \gtrsim 2$) Red Galaxies (DRGs) by selecting galaxies with $J - K_s > 2.3$ (Franx et al. 2003). At similar restframe optical luminosities, DRGs typically have older stellar ages (1-3 Gyr), larger stellar masses ($\sim 10^{11} M_\odot$), more dust ($A_V = 1-3 \text{ mag}$) and higher star formation rates ($\text{SFR} \sim 100 M_\odot \text{ yr}^{-1}$) than the complementary population of high redshift galaxies selected using the Lyman break technique (Steidel et al. 1999; Shapley et al. 2001, 2003; Förster-Schreiber et al. 2004; van Dokkum et al. 2004). The observed properties of LBGs suggests that they are on average younger than the more evolved DRGs.

Morphological studies of Lyman Break Galaxies (LBGs) using the Hubble Space Telescope (HST) suggest that they are compact ($r_e \sim 2 \text{ kpc}$) and have very similar restframe UV and optical morphologies (Giavalisco et al. 1996; Lowenthal et al. 1997; Dickinson 2000; Ferguson et al. 2004; Bouwens et al. 2004). Here we present the first

space-based study of the restframe optical morphologies of DRGs. We take advantage of the Hubble Ultra Deep Field (UDF) NICMOS and ACS imaging of a $2.4 \times 2.4 \text{ arcmin}^2$ area of the sky in the Chandra Deep Field South (CDFs), to study restframe UV to optical morphologies of DRGs. The unprecedented depth⁶ of the data allow for detailed morphological studies, even of very low surface brightness components.

In the following we assume a standard cosmology with $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, and $h = 0.7$ (Spergel et al. 2003). We refer to the HST filters F435W, F606W, F775W, F850LP, F110W and F160W as B_{435} , V_{606} , i_{775} , z_{850} , J_{110} and H_{160} . Magnitudes are in the Vega system, unless otherwise noted.

2. SELECTION

In addition to the public UDF data (Beckwith et al. 2004; Thompson et al. 2004) we use a photometric catalog consisting of ACS data (in the B_{435} , V_{606} , i_{775} and z_{850} -bands) taken as part of the GOODS survey,

¹ Based on observations made with the NASA/ESA *Hubble Space Telescope*, which is operated by the Association of Universities for Research in Astronomy, Inc, under NASA contract NAS5-26555

² Department of Astronomy, Yale University, New Haven, CT 06520-8101, email: toft@astro.yale.edu, dokkum@astro.yale.edu

³ Leiden Observatory, Postbus 9513, 2300 RA Leiden, Netherlands, email: franx@strw.leidenuniv.nl, mariska@strw.leidenuniv.nl

⁴ Steward Observatory, University of Arizona, 933 Cherry Avenue, Tucson, AZ 85721, email: thompson@cosmos.as.arizona.edu

⁵ University of California at Santa Cruz, 1156 High Street, Santa Cruz, CA 95064, email: gdi@ucolick.org, bouwens@ucolick.org

⁶ The 10σ AB limits in the filters F435W, F606W, F775W, F850LP, F110W, F160W are 29.2, 30.0, 29.7, 28.7, 27.0, 27.0., (Beckwith et al. 2004; Thompson et al. 2004)




FIG. 1.— Gallery of the six galaxies in the UDF with $K_s < 21.5$ and $J - K_s > 2.3$. The filters shown are (from left to right): $B_{435} + V_{606}$, $i_{775} + z_{850}$, J_{110} , H_{160} . The size of the stamps is $3''.6 \times 3''.6$. North is up, east is to the left. The images have been normalized such that a galaxy with constant f_λ would appear the same in all images. The galaxies appear regular in the NICMOS images, and show a range of sizes.

combined with deep VLT/ISAAC data (in the J , H and K_s -bands) taken as part of the ESO Imaging Survey (EIS) (van der Wel et al. 2004). In the following we refer to this catalog as the GOODS/EIS catalog.

We use the GOODS/EIS catalog rather than the UDF catalog for selecting galaxies as it includes the same J and K_s filters that have been used previously (Franx et al. 2003). The additional depth of the UDF J_{110} -band observation (relative to the ISAAC J -band observations) is less important as we are concentrating on relatively bright galaxies. Galaxies were selected using the criteria: $K_s < 21.5$, $J - K_s > 2.3$. Six galaxies in the UDF match these criteria. A postage stamp gallery of the sample is shown in Fig. 1.

3. PHOTOMETRY AND PHOTOMETRIC REDSHIFTS

Photometric redshifts were derived both from $1''.8$ diameter aperture photometry derived from Point Spread Function (PSF)-matched ACS/NICMOS UDF images, and from $2''$ diameter aperture photometry available in the GOODS/EIS catalog. The GOODS/EIS dataset was

found to provide stronger constraints on the redshifts than the UDF data set, due to the inclusion of the K_s -band and the fact that the ISAAC J -band is redder and narrower than the NICMOS J_{110} band. Hyperz (Bolzonella et al. 2000) was used to derive photometric redshifts. To limit the degrees of freedom in the fits, we used four empirical templates (E, Sbc, Scd and Im) to derive the best fitting photometric redshifts. In Tab 1, we list the best fitting photometric redshifts and other properties of the sample. The median photometric redshift of the sample is 2.1 ± 0.7 , in good agreement with the median photometric redshift found for DRGs in the HDFS (Franx et al. 2003). One of the six galaxies in the sample has a photometric redshift significantly lower than 2. The surface density of $z_{phot} \gtrsim 2$ DRGs to $K_s = 21.5$ is $\simeq 0.9 \text{ arcmin}^{-2}$, consistent with $\simeq 1.0 \text{ arcmin}^{-2}$ to $K_s = 21.7$ found for a much larger sample in the MS1054 field (Förster-Schreiber et al. 2004). Following Franx et al. (2003) and Förster-Schreiber et al. (2004), we next fixed the redshifts and fitted three Bruzual & Charlot (2003) models (BC): a constant star formation model (CSF) with dust, a “Tau” model (exponentially declining star formation with a characteristic timescale of $\tau = 1 \text{ Gyr}$) with dust, and a Single instantaneous burst model without dust. In Fig 2. we show SEDs of the six galaxies in the sample, along with the best fitting redshifted BC models (full black curves). It is interesting to note that all six galaxies are better fitted by the “Tau” model with ongoing star formation and significant amounts of dust, than the single burst model with no dust (shown in dashed red curves).

4. MORPHOLOGIES

The DRGs in our sample show a range of morphologies: in the NIR, three are extended and three are compact. Of the galaxies with $z_{phot} \gtrsim 2$, three are compact and two are extended. The observed NIR (restframe optical) morphology of the $z_{phot} \gtrsim 2$ galaxies look quite different from the observed optical (restframe UV) morphology, highlighting the importance of deep NIR imaging of high redshift galaxies. In the NIR, all the $z_{phot} \gtrsim 2$ galaxies have red central components which dominate the NICMOS H_{160} -band images, while distinct blue off-center features show up in (and often dominate) the ACS images. We verified that these differences were not due to the different PSF and noise properties of the ACS and NICMOS images.

The most pronounced differences between ACS and NICMOS are seen in two galaxies: UDF-DRG1, which has a luminous red bulge which dominates the light in the NIR bands and a more extended, patchy blue disk which dominates the light in the optical bands; and UDF-DRG6, which in the NIR bands is a single compact source, but in the optical bands is composed of four close compact sources of comparable brightness. From the optical bands alone these two galaxies could easily have been classified as highly irregular galaxies, or galaxies undergoing major merging, while their NIR morphologies reveal that they are more likely relatively relaxed galaxies with patches of ongoing star formation.

The extraordinary depth of the UDF data allow us to test whether the blue “blobs” seen in UDF-DRG 1, 3, 5 and 6 are due to substructure in the




FIG. 2.— SEDs (f_λ versus λ) of the six galaxies in the sample. Solid circles are photometry from the GOODS/EIS catalog (normalized by f_λ in Ks -band). The full black curves are the best fitting (SB, Tau or CSF) BC models and the red dashed curves are the best fitting SB models (see text for details). The Tau model (with some dust extinction) provides the best fit to all six galaxies. The CSF models provide almost as good fits, but require more dust. The formal errors on the redshift from HYPERZ are in the range ~ 0.1 - 0.2 , as are error estimates from monte carlo simulations.

galaxies, or chance alignment with unrelated sources at different redshifts. To do this we placed small (3 pix) apertures on the positions of the individual blobs (indicated on the $B_{435} + V_{606}$ images in Fig. 1) in the PSF-matched images and derived their SEDs and photometric redshift. These are all consistent with the blobs being substructure at the redshift of the main galaxies, although we note that the photometry in neighboring blobs are not fully independent.

5. SIZES

To better quantify the differences and similarities of the galaxies and their appearance at different wavelengths we modeled their surface brightness distributions using GALFIT (Peng et al. 2002). We fitted a Sersic profile (convolved with the PSF, estimated from a calibration star): $I(r) = I(0)\exp[b_n(r/r_e)^{1/n}]$ which has been shown to be a good representation of the surface brightness of a range of galaxy types from ellipticals ($n = 4$) to disks ($n = 1$). Leaving the radial shape parameter n and the effective radius r_e free in the fitting resulted in a mean value of $n = 1.4 \pm 0.9$. Since there is some degeneracy between n and r_e in the fits, we next fixed n to 1, to be able to compare the derived sizes of the galaxies with each other and with other high redshift galaxies. The resulting sizes (listed in Tab. 1) range from $0''.1$ to $0''.5$. At NIR wavelengths (in the H_{160} -band), the mean measured circularized radius⁷ of the five $z_{phot} \gtrsim 2$ galaxies is $\langle r_e \rangle_{NIR} = 0''.23 \pm 0''.16$. At the redshifts of the galaxies this corresponds to a mean physical size of 1.9 ± 1.4 kpc. At optical wavelengths most of the sizes are slightly larger, but less well defined due to the patchy light distribution.

⁷ defined as $r_{e,c} = r_e \sqrt{(1 + \epsilon)}$, where r_e is the effective radius encompassing half the flux, measured along the semi-major axis, and $\epsilon = 1 - a/b$ is the ellipticity of light distribution




FIG. 3.— Comparison of red galaxies selected here from their $J - Ks$ color (large solid symbols), from their $I_{775} - H_{160}$ color and z_{phot} (large open symbols, Chen & Marzke 2004), and from their $z_{850} - 3.6\mu m$ ($IRAC$) color (small filled (pink) symbols, Yan et al. 2004). The colors and magnitudes of the galaxies are from the GOODS/EIS catalog. Four of the Chen & Marzke (2004) and four of the Yan et al. (2004) galaxies are too faint to be detected in our Ks -band data. The three samples only have two galaxies in common (UDF-DRG4 and 6)

All the DRGs are resolved in the ACS images. To determine whether the “compact” galaxies (UDF-DRG 3, 4 and 6) are resolved in the NICMOS data we repeated the size-fitting analysis on three stars (two stars in the field and a calibration star), to derive the mean size of “bonafide” unresolved objects: $\langle r_e \rangle_{stars} = 0''.06 \pm 0''.03$. From this we conclude that UDF-DRG4 is marginally resolved, while UDF-DRG3 and 6 are consistent with being unresolved. Varying the fitting regions, and using different stars to model the PSF result in little variation in the derived r_e .

The derived sizes can be compared to those of $z \sim 3$ LBGs. Ferguson et al. (2004) find a mean size $\langle r_e \rangle \sim 2.3$ kpc, and Bouwens et al. (2004) find $\langle r_e \rangle \sim 1.8$ kpc. As the luminosity range of the $z_{phot} \gtrsim 2$ galaxies in our sample ($M_V = -21.4$ to -23.3) is comparable to that of the LBG samples we can compare the sizes directly (see Trujillo et al. 2004). The mean size for $z_{phot} \gtrsim 2$ DRGs derived here is similar within the errors to the mean size of LBGs.

A possible caveat of this comparison is the fact that we are comparing DRG sizes derived in the restframe optical, with LBG sizes derived in the restframe UV. However, the difference between restframe UV and optical sizes derived here for our small sample of DRGs (see Tab.1) are much smaller than the galaxy-galaxy variation in sizes.

6. AGN CONTRIBUTION

Two of the DRGs in our sample are detected in the 1Ms Chandra Deep Field South X-ray catalog (Giacconi et al. 2002) to $f_x > 10^{-16}$ ergs $\text{cm}^{-2}\text{s}^{-1}$. The $[0.5-2.0]$ keV band X-ray fluxes of UDF-DRG 1 and 3 ($f_x = 73$ and 81×10^{-17} ergs $\text{cm}^{-2}\text{s}^{-1}$ respectively) are comparable to

TABLE 1
PROPERTIES OF DISTANT RED GALAXIES IN THE UDF

Id ^a	K_s	$J - K_s$	z_{phot}^b	$r_{e,c}^c$
UDF-DRG1 /41	19.38	2.42	2.2	0''35/0''40
UDF-DRG2 /372	20.69	2.32	1.2	0''17/0''19
UDF-DRG3 /290	20.70	2.66	1.9	0''09 ^d /0''03
UDF-DRG4 /746	20.93	3.26	3.4	0''14/0''12
UDF-DRG5 /899	19.81	2.45	1.8	0''49/0''53
UDF-DRG6 /1207	21.00	3.24	3.4	0''08 ^d /0''10

^a Id in this letter / Id in the public NICMOS UDF catalog (vers.2.0)

^b Photometric redshift derived from the GOODS/EIS catalog

^c Circularized effective radius derived in the H_{160}/V_{606} filters. The typical error is $\sim 0''.1$

^d Consistent with unresolved

the fluxes of the two DRGs with AGNs in the MS1054-03 field (Rubin et al. 2004). The corresponding X-ray luminosities are too bright to be supported by star formation and must thus be of AGN origin (the X-ray flux, due to star formation, expected from DRGs $f_x \simeq 10^{-17}$ ergs $s^{-1}cm^{-2}$ (Rubin et al. 2004) is below the detection limit). The fact that two of the $z_{phot} \gtrsim 2$ galaxies certainly hosts AGN allow us to test the AGNs contribution to the rest-frame optical emission. UDF-DRG1 is clearly resolved in the NIR bands, so its red restframe optical color is probably due to evolved stars, rather than the AGN. UDF-DRG3 on the other hand is unresolved in the NIR bands, and could thus be dominated by the AGN, rather than stars, in its restframe optical emission.

7. DISCUSSION

Two other studies have recently identified red galaxies in the UDF. Chen & Marzke (2004) select 9 galaxies with $I_{775} - H_{160} > 2$ and $z_{phot} > 2.5$, six of which are argued to be best described by evolved templates with no dust, and three by dusty starbursts. Yan et al. (2004) combine the HST data with mid-infrared observations obtained with the Infra Red Array Camera (IRAC) onboard the Spitzer Space Telescope, to select 10 galaxies with $f_\nu(3.6\mu m)/f_\nu(z_{850}) > 20$, and argue that their SEDs are well represented by a two component model consisting of an old evolved population and a secondary young population. In Fig. 3 we plot $J - K_s$ vs K_s for the three samples of red galaxies in the UDF.

The two reddest DRGs in our sample make it into both the Chen & Marzke (2004) sample (Object 09024 and 01927 in their Table 1) and the Yan et al. (2004) sam-

ple (object 8 and 9 in their Table 1). Of the remaining seven/eight galaxies in the Chen & Marzke (2004)/Yan et al. (2004) samples, four/four are too faint to be detected in our data, while three/four are either too blue or too faint to make our selection criterion. We conclude that while the different selection methods show some overlap, they are not redundant.

We note, that while the photo- z 's derived for the two galaxies present in all three samples agree reasonably well (± 0.2), the model parameters are uncertain: we find both galaxies to be best fit by a 1Gyr Tau model with dust; Chen & Marzke (2004) find UDF-DRG4 to be best fit by a 0.3Gyr Tau model without dust, and UDF-DRG6 by a dusty CSF model; and Yan et al. (2004) fits a two component model without dust to both galaxies. This demonstrates that while photometric redshifts derived from fits to broadband SEDs are relatively robust, parameters of the fitted models are in general less well determined and should be interpreted with caution.

The most characteristic property of the restframe optical and UV morphologies of the DRGs studied here is their diversity. Some are extended, some are compact, some have very different restframe optical and UV morphologies, while some look similar. The conclusion that can be drawn is that most of the DRGs in the present sample have composite stellar populations, with central populations of red evolved stars dominating the restframe optical morphologies and younger more patchy distributed blue stars dominating the restframe UV morphologies, while LBGs have similar restframe optical and UV morphologies (Dickinson 2000), and thus may be uniformly younger systems with star forming regions so luminous that they dominate even the restframe optical light. These results are in agreement with those derived from spectra and broadband SEDs of DRGs (Förster-Schreiber et al. 2004) and LBGs (Pettini et al. 2001; Shapley et al. 2003). There might, however, be some overlap in the properties of restframe optical and UV selected galaxies as a small fraction of restframe UV selected LBGs at $z \simeq 3$ and Bm/Bx galaxies at $z \simeq 2$ have recently been shown to have red optical colors and ages similar to the DRGs (Steidel et al. 2004; Shapley et al. 2004).

Support from the Danish Natural Science Research Council, and from NASA through HST grant GO-09803-10-A and LTSA NNG04GE12G is gratefully acknowledged.

REFERENCES

- Beckwith et al. 2004, in prep.
 Bolzonella, M., Miralles, J.-M., & Pell, R. 2000, *A&A*, 363, 476
 Bouwens, R. J., Illingworth, G. D., Blakeslee, J. P., Broadhurst, T. J., & Franx, M. 2004, *ApJ*, 611, L1
 Bruzual, A. G. & Charlot, S. 2003, *Priv.comm*
 Chen, H., & Marzke, R. O. 2004, *ApJ*, 615, 603
 Daddi, E., Röttgering, H. J. A., Labbé, I., et al. 2003, *ApJ*, 588, 50
 Dickinson, M. 2000, *RSPA*, 358, 2001
 Förster-Schreiber, N. M., van Dokkum, P. G., Franx, M., et al. 2004, *ApJ*, 616, 40
 Ferguson, H. C., Dickinson, M., Giavalisco, M., et al. 2004, *ApJ*, 600, L107
 Franx, M., Labbé, I., Rudnick, G., et al. 2003, *ApJ*, 587, L79
 Giacomini, R., Zirm, A., Wang, J., et al. 2002, *ApJS*, 139, 369
 Giavalisco, M., Steidel, C. C., & Macchetto, F. D. 1996, *ApJ*, 470, 189
 Labbé, I., Franx, M., Rudnick, G., et al. 2003, *AJ*, 125, 1107
 Lowenthal, J. D., Koo, D. C., Guzman, R., et al. 1997, *ApJ*, 481, 673
 Peng, C. Y., Ho, L. C., Impey, C. D., & Rix, H. 2002, *AJ*, 124, 266
 Pettini, M., Shapley, A. E., Steidel, C. C., Cuby, J., Dickinson, M., Moorwood, A. F. M., Adelberger, K. L., & Giavalisco, M. 2001, *ApJ*, 554, 981
 Rubin, K., van Dokkum, P., Coppi, C., et al. 2004, *ApJ*, 613, L5
 Shapley, A. E., Steidel, C. C., Adelberger, K. L., et al. 2001, *ApJ*, 562, 95
 Shapley, A. E., Steidel, C. C., Pettini, M., & Adelberger, K. L. 2003, *ApJ*, 588, 65
 Shapley, A. E., Erb, D. K., Pettini, M., Steidel, C. C., & Adelberger, K. L. 2004, *ApJ*, 612, 108
 Spergel, D. N., Verde, L., Peiris, H. V., et al. 2003, *ApJS*, 148, 175
 Steidel, C. C., Adelberger, K. L., Giavalisco, M., Dickinson, M., & Pettini, M. 1999, *ApJ*, 519, 1
 Steidel, C. C., Shapley, A. E., Pettini, M., Adelberger, K. L., Erb, D. K., Reddy, N. A., & Hunt, M. P. 2004, *ApJ*, 604, 534
 Thompson, R., Illingworth, G., Bouwens, R., et al. 2004, *AJ* (submitted)

Trujillo, I., et al. 2004, ApJ, 604, 521
van der Wel et al. 2004, in prep
van Dokkum, P. G., et al. 2003, ApJ, 587, L83

van Dokkum, P. G., et al. 2004, ApJ, 611, 703
Yan, H., et al. 2004, ApJ, 616, 63